

Topology and its Applications 48 (1992) 179–184
North-Holland

179

The quasitopos hull of the category of uniform spaces – a correction

Jiří Adámek

Faculty of Electrical Engineering, Technical University of Prague, Technická 2, Prague 6, Czechoslovakia

Jan Reiterman

Faculty of Nuclear Science and Technical Engineering, Technical University of Prague, Trojanova 13, Prague 2, Czechoslovakia

Received 27 October 1989

Revised 2 March 1991 and 29 August 1991

Abstract

Adámek, J. and J. Reiterman, The quasitopos hull of the category of uniform spaces – a correction, *Topology and its Applications* 48 (1992) 179–184.

We present a description of the quasitopos hull of the category of uniform spaces. This corrects our earlier description published in *Topology Appl.* 27.

Keywords: Concrete quasitopos, category of uniform spaces, quasitopos hull, semi-uniformity.

AMS (MOS) Subj. Class.: 54B30, 18B25, 18B30, 18F10, 18B15.

In our paper [4] we attempted to describe the quasitopos hull of the category **Unif** of uniform spaces. Unfortunately, that description was incorrect, and the reason was a mistake in [2] where it was claimed that, given a topological category \mathcal{K} which is a full, finally dense subcategory of a quasitopos \mathcal{Q} then the initial lifts of all $[K_1, K_2]^\#$ with $K_1, K_2 \in \mathcal{K}$ form the quasitopos hull of \mathcal{K} . (We denote power objects by $[A, B]$ and objects representing strong partial morphisms with codomain A by $A^\#$.) The statement has been corrected in [5] as follows:

Correspondence to: Dr. J. Adámek, Technical University of Prague, FEL CVUT, Suchbátarova 2, Praha 6, Czechoslovakia.

Theorem. Let \mathcal{K} be a topological category which is a full, finally dense subcategory of a concrete quasitopos \mathcal{Q} . Then \mathcal{Q} is a quasitopos hull of \mathcal{K} iff all objects $[K_1, K_2^\#]$ with $K_1, K_2 \in \mathcal{K}$ are initially dense in \mathcal{Q} .

Corollary. Let \mathcal{K} be a topological category which is a full, finally dense subcategory of a concrete quasitopos \mathcal{Q} . Then the initial hull of all objects $[K_1, K_2^\#]$ of \mathcal{Q} with $K_1 \in \mathcal{Q}$ and $K_2 \in \mathcal{K}$ forms a quasitopos hull of \mathcal{K} .

In fact, from the proof of the above theorem it follows readily that the quasitopos hull of \mathcal{K} is the initial hull of all objects $[K_1, K_2^\#]$ with $K_1, K_2 \in \mathcal{K}$. For $K_1 \in \mathcal{Q} - \mathcal{K}$ use a final sink $(A_i \xrightarrow{\alpha_i} K_1)_{i \in I}$ with $A_i \in \mathcal{K}$: then $([K_1, K_2^\#] \xrightarrow{[\alpha_i, 1]} [A_i, K_2^\#])_{i \in I}$ is an initial source, thus, $[K_1, K_2^\#]$ lies in the quasitopos hull.

In what follows we work with *semi-uniform spaces*, see [11], i.e., pairs (X, α) where α is a filter (w.r.t. the ordering by inclusion) of symmetric vicinities of the diagonal in $X \times X$ (i.e., of sets $U \subseteq X \times X$ with $\Delta_X \subseteq U = U^{-1}$). Let **SUnif** be the category of semi-uniform spaces, where morphisms $f: (X, \alpha) \rightarrow (Y, \beta)$ are functions $f: X \rightarrow Y$ such that $U \in \beta$ implies $(f \times f)^{-1}(U) \in \alpha$.

Unif is, obviously, a full subcategory of **SUnif**. It is finally dense in **SUnif** because, as proved in [7], every semi-uniform space is a quotient of a uniform space. It is easy to verify that both quotients and finite coproducts are stable under pullbacks in **SUnif**. Thus, by [4, Remark 8], we have the concrete quasitopos $A^*\mathbf{SUnif}$ of *filters of semi-uniformities*: objects are pairs (X, \mathcal{A}) where X is a set and \mathcal{A} is a filter of semi-uniform spaces (Y, β) , $Y \subseteq X$, under the ordering $<$ defined by $(Y, \beta) < (Y', \beta')$ iff $Y \subseteq Y'$ and the inclusion is a morphism $(Y, \beta) \rightarrow (Y', \beta')$ of **SUnif**. *Filter* means that

- (a) $(Y, \beta) \in \mathcal{A}$ implies $(Y', \beta') \in \mathcal{A}$ whenever $(Y', \beta') < (Y, \beta)$,
- (b) for $(Y', \beta'), (Y'', \beta'') \in \mathcal{A}$ there exists $(Y, \beta) \in \mathcal{A}$ with $(Y', \beta') < (Y, \beta)$ and $(Y'', \beta'') < (Y, \beta)$,
- (c) for each $x \in X$ there exists $(Y, \beta) \in \mathcal{A}$ with $x \in Y$.

Morphisms $f: (X, \mathcal{A}) \rightarrow (X', \mathcal{A}')$ are functions $f: X \rightarrow X'$ such that for each $(Y, \beta) \in \mathcal{A}$ there exists $(Y', \beta') \in \mathcal{A}'$ with $f(Y) \subseteq Y'$ and such that the domain-range restriction of f defines a morphism $(Y, \beta) \rightarrow (Y', \beta')$ of **SUnif**.

We consider **SUnif** as a full, concrete subcategory of $A^*\mathbf{SUnif}$ by identifying each $(X, \alpha) \in \mathbf{SUnif}$ with $(X, \{(Y, \beta): (Y, \beta) < (X, \alpha)\})$.

Definition. (1) A semi-uniformity α is called *simple* if it has a base formed by a single vicinity U , i.e., α consists of all symmetric supersets of U . The semi-uniformity is then denoted by $[U]$.

(2) A filter of semi-uniformities is called *simple* if it has a basis of simple semi-uniformities. That is, (X, \mathcal{A}) is simple if for each $(Y, \beta) \in \mathcal{A}$ there is $(Y', [U]) \in \mathcal{A}$ with $(Y, \beta) < (Y', [U])$.

(3) A filter of semi-uniformities is called *saturated* if it is an intersection of simple filters.

(4) The full subcategory of $A^*\mathbf{SUnif}$ of all saturated filters is denoted by \mathbf{SF} .

Lemma 1. *The power object of semi-uniform spaces $A = (X, \alpha)$, $B = (Y, \beta)$ in the category $A^*\mathbf{SUnif}$ is the following saturated filter of semi-uniformities:*

$$[A, B] = (\text{hom}(A, B), \mathcal{C})$$

where \mathcal{C} consists of all semi-uniform spaces (M, μ) satisfying the following conditions:

- (1) $M \subseteq \text{hom}(A, B)$ is an equicontinuous set, i.e., for each $V \in \beta$ there exists $U \in \alpha$ with $\bigcup_{f \in M} f \times f(U) \subseteq V$; in other words, there is a function $r: \beta \rightarrow \alpha$ coherent in the sense that $\bigcup_{f \in M} f \times f(rV) \subseteq V$ for all $V \in \beta$.
- (2) μ contains, for some coherent function $r: \beta \rightarrow \alpha$, all the vicinities $V^{(r)} = \{(f, g) \in M \times M: f \times g(rV) \subseteq V\}$ for $V \in \beta$, i.e., $\{V^{(r)}: V \in \beta\} \subseteq \mu$.

Proof. (I) Evaluation is a morphism $\text{ev}: [A, B] \times A \rightarrow B$ in $A^*\mathbf{SUnif}$. To prove this, it clearly suffices to show, for each equicontinuous set $M \subseteq \text{hom}(A, B)$ and every coherent function r , that the restriction of the evaluation is a morphism $\text{ev}: (M, \mu) \times A \rightarrow B$ of \mathbf{SUnif} , where μ is the semi-uniformity with the subbasis $\{V^{(r)}: V \in \beta\}$. This is obvious: for each $V \in \beta$ we have vicinities $V^{(r)}$ and $V^* = rV \in \alpha$ such that given $(f, g) \in V^{(r)}$ and $(x, y) \in V^*$ then

$$(f(x), g(y)) \in (f \times g)(V^*) \subseteq V.$$

(II) Given an object D in $A^*\mathbf{SUnif}$ and a morphism $f: D \times A \rightarrow B$, the function $\hat{f}: D \rightarrow [A, B]$, $d \mapsto f(d, -)$, is a morphism, too. We prove this first in case $D = (T, \delta)$ is a semi-uniform space:

- (a) Each $f(d, -): A \rightarrow B$ is clearly uniformly continuous.
- (b) \hat{f} is a morphism, i.e., $T = \hat{f}(M)$ is an equicontinuous set and, for some coherent map r , the range restriction of \hat{f} is uniformly continuous from D to (M, μ) . In fact, for each $V \in \beta$ there exist $sV \in \delta$ and $rV \in \alpha$ such that

$$(d_1, d_2) \in sV \text{ and } (x, y) \in rV \text{ imply } (f(d_1, x), f(d_2, y)) \in V.$$

It follows that the map $r: \beta \rightarrow \alpha$ is coherent: for each $f(d, -) \in M$ we have $(d, d) \in sV$ and thus $(f(d, -) \times f(d, -))(rV) \subseteq V$. The continuity of the restriction of \hat{f} is obvious: for each $V^{(r)}$ where $V \in \beta$ we have $sV \in \delta$ for which $(d_1, d_2) \in sV$ implies $(\hat{f}(d_1), \hat{f}(d_2)) \in V^{(r)}$.

Next, let D be a filter of semi-uniformities D_i , $i \in I$. Since each of the restrictions $f_i: D_i \times A \rightarrow B$ of $f: D \times A \rightarrow B$ is a morphism, it follows from the above that each $\hat{f}_i: D_i \rightarrow [A, B]$ is a morphism and thus $\hat{f}: D \rightarrow [A, B]$ is a morphism since the sink of inclusion maps $(D_i \hookrightarrow D)_i$ is final in $A^*\mathbf{SUnif}$.

(III) Finally, let us verify that $[A, B]$ is a saturated filter. If B is a simple semi-uniform space, then $[A, B]$ is, obviously, a simple filter of semi-uniformities. In general, each semi-uniform space B has an initial source $(B \xrightarrow{\text{id}} B_i)_i$ with each B_i simple, and this yields an initial source $([A, B] \xrightarrow{[A, \text{id}]} [A, B_i])_i$ whose codomains are simple filters—thus the domain is saturated. \square

Lemma 2. *For each semi-uniform space K the object $K^\#$ of $A^*\mathbf{SUnif}$ is a semi-uniform space.*

Proof. It is easy to verify that if $K = (X, \alpha)$ then $K^\# = (X \cup \{\infty\}, \alpha^\#)$ where $\alpha^\# = \{U \cup (\{\infty\} \times X) \cup (X \times \{\infty\}) \cup \{\infty, \infty\} : U \in \alpha\}$. \square

Characterization Theorem. *The category \mathbf{SF} of saturated filters of semi-uniformities is a quasitopos hull of \mathbf{Unif} .*

Proof. \mathbf{SUnif} is clearly closed under quotients in $A^*\mathbf{SUnif}$. Since each semi-uniform space is a quotient of a uniform space (in \mathbf{SUnif}), it follows that not only \mathbf{SUnif} but even \mathbf{Unif} is finally dense in $A^*\mathbf{SUnif}$. By the above corollary, it is sufficient to prove that the initial hull of all power objects $[K_1, K_2^\#]$ where K_2 is a uniform space is equal to \mathbf{SF} . By Lemmas 1 and 2, these power objects lie in \mathbf{SF} for $K_1 \in \mathbf{SUnif}$. For K_1 general choose a final sink $(L_i \rightarrow K_1)$ with $L_i \in \mathbf{Unif}$; we obtain an initial lift $([K_1, K_2^\#] \rightarrow [L_i, K_2^\#])$ in $A^*\mathbf{SUnif}$. Since \mathbf{SF} is clearly closed in $A^*\mathbf{SUnif}$ under initial lifts, it follows that $[K_1, K_2^\#] \in \mathbf{SF}$.

Thus, it is sufficient to prove that each simple filter (X, \mathcal{A}) of semi-uniformities is an initial lift of an object $[P, R]$ where P is a semi-uniform space and $R = S^\#$ for some uniform space S . Let $R = \{0, 1\}^\#$ where $\{0, 1\}$ is the discrete uniform space, i.e., R is the simple semi-uniform space on the set $\{0, 1, \infty\}$ whose generating vicinity is

$$W = \{0, 1, \infty\} \times \{0, 1, \infty\} - \{(0, 1), (1, 0)\}.$$

Let \mathcal{A}_0 be the basis of \mathcal{A} formed by all simple semi-uniform spaces in \mathcal{A} , and for each $A \in \mathcal{A}_0$, denote by V_A the generating vicinity. Denote further by P the following semi-uniform space: its underlying set is $X \times X \times \mathcal{A}_0$ and the following vicinities $U_A, A \in \mathcal{A}_0$, form a basis of its semi-uniformity:

$$U_A = \{(r, s) \in P \times P : (x, y) \in V_A \Rightarrow ((e(x)(r), e(y)(s)) \in W)\}$$

where $e : X \rightarrow R^P$ is defined by

$$e(x)(r) = \begin{cases} 0, & \text{if } r = (x, x, A) \text{ for some } A \in \mathcal{A}_0, \\ 1, & \text{if } r = (x, y, A) \text{ for some } A \in \mathcal{A}_0 \\ & \text{with } (x, y) \notin V_A \text{ and } y \in A, \\ \infty, & \text{else.} \end{cases}$$

(To verify that $\{U_A\}$ is a basis of vicinities observe that each U_A contains the diagonal of P , and that $U_A \subseteq U_{A_1} \cap U_{A_2}$ holds whenever $V_{A_1} \cup V_{A_2} \subseteq V_A$.) We are going to prove that

$$e : (X, \mathcal{A}) \rightarrow [P, R] \text{ is initial in } A^*\mathbf{SUnif}.$$

(a) For each $x \in X$, $e(x) : P \rightarrow R$ is uniformly continuous since $e(x) \times e(x)(U_{\{x\}}) \subseteq W$.

(b) e is a morphism since for each $A \in \mathcal{A}_0$, $M = e(A)$ is equicontinuous, and the (coherent) function r given by $rW = U_A$ has the property that the domain-range restriction of e is uniformly continuous from A to $(M, \{V: V \text{ vicinity}, W^{(r)} \subseteq V\})$. In fact, if $(x, y) \in V_A$ then

$$(e(x), e(y)) \in W^{(r)} = \{(f, g) \in M \times M: f \times g(U_A) \subseteq W\}.$$

(c) To show that e is initial, suppose that $D = (T, \delta)$ is a semi-uniform space with $T \subseteq X$ such that the domain-range restriction of e is a morphism $e': D \rightarrow [P, R]$ (i.e., there exists an equicontinuous set $M \subseteq \text{hom}(P, R)$ and a coherent function r [a choice of a vicinity rW in P] such that $e': D \rightarrow (M, \{V: W^{(r)} \subseteq V\})$ is uniformly continuous). We are going to prove that then $D < A$ for some $A \in \mathcal{A}$. The uniform continuity of e' means that there exists a vicinity $V^* \in \delta$ such that $(e \times e)(V^*) \subseteq W^{(r)}$. Since rW is a vicinity in P , it contains some U_A ($A \in \mathcal{A}_0$), and thus

$$(x, y) \in V^* \text{ implies } (e(x) \times e(y))(U_A) \subseteq W.$$

We are going to prove that $D < A$.

(a) $T \subseteq A$. If contrary, choose $t \in T - A$; further, choose $a \in A$. Since $e(t)(t, t, A) = 0$ and $e(t)(t, a, A) = 1$, we have, for $r = (t, t, A)$ and $s = (t, a, A)$:

$$(e(t)(r), e(t)(s)) \notin W.$$

This is a contradiction since, clearly, $(t, t) \in V^*$ and $(r, s) \in U_A$.

(b) To prove $D < A$, it is sufficient to verify that $V^* \subseteq V_A$. If contrary, choose $(x, y) \in V^* - V_A$ and put $r = (x, x, A)$ and $s = (y, x, A)$. Then $(r, s) \in U_A$ but $(e(x)(r), e(y)(s)) = (0, 1) \notin W$, a contradiction.

This concludes the proof that $e: D \rightarrow [P, R]$ is an initial morphism. \square

Example. \mathbf{SF} is a proper subcategory of $A^*\mathbf{SUnif}$; one can prove that the following filter (X, \mathcal{B}) is not saturated: Let X be an infinite set and let a subbase of \mathcal{B} be formed by all semi-uniform spaces $(X, \alpha_{\mathcal{F}})$ where \mathcal{F} is a free ultrafilter on $X \times X - \Delta_X$ and

$$\alpha_{\mathcal{F}} = \{U \cup \Delta_X: U = U^{-1} \in \mathcal{F}\}.$$

Remark. In [3] we have described the cartesian closed topological hull of the category of uniform spaces as the category of bornological uniform spaces (X, α, \mathcal{U}) where α is a uniformity on X and \mathcal{U} is a bornology on X naturally related to each other.

In contrast to bornological uniform spaces, in the concept of saturated filter of semi-uniformities, (1) there is not a single semi-uniformity but various semi-uniformities on various subsets and (2) the bornology (of all underlying sets of the members of the filter) is not related to the structure of semi-uniformities.

Acknowledgement

We are grateful to Oswald Wyler for pointing out some mistakes in an earlier version of our paper.

References

- [1] J. Adámek, *Theory of Mathematical Structures* (Reidel, Dordrecht, 1983).
- [2] J. Adámek, Classification of concrete categories, *Houston J. Math.* 12 (1986) 305–326.
- [3] J. Adámek and J. Reiterman, Cartesian closed hull of the category of uniform spaces, *Topology Appl.* 19 (1985) 261–276.
- [4] J. Adámek and J. Reiterman, The quasitopos hull of the category of uniform spaces, *Topology Appl.* 27 (1987) 97–104.
- [5] J. Adámek, J. Reiterman and F. Schwarz, On universally topological hulls and quasitopos hulls, *Houston J. Math.* 17 (1991) 375–383.
- [6] Ph. Antoine, Etude élémentaire des catégories d'ensembles structures I, II, *Bull. Soc. Math. Belg.* 18 (1966) 142–164; *ibid.* 18 (1966) 387–414.
- [7] E. Čech, *Topological Spaces* (Academia, Prague, 1966).
- [8] H. Herrlich, Hereditary topological constructs, in: *Proceedings of the 6th Prague Topological Symposium* (Heldermann, Berlin, 1988) 249–262.
- [9] H. Herrlich, On the representability of partial morphisms in **Top** and in related constructs, *Lecture Notes in Mathematics* 1348 (Springer, Berlin, 1988) 144–153.
- [10] H. Herrlich and L.D. Nel, Cartesian closed topological hulls, *Proc. Amer. Math. Soc.* 62 (1977) 215–222.
- [11] J.R. Isbell, *Uniform Spaces* (Amer. Math. Soc., Providence, RI, 1964).